

# Generalization and simplifications of generating functions for Jacobi, Gegenbauer, Chebyshev and Legendre polynomials with definite integrals

Howard S. Cohl <sup>†</sup> and Connor MacKenzie <sup>‡†</sup>

<sup>†</sup> *Applied and Computational Mathematics Division, Information Technology Laboratory,  
National Institute of Standards and Technology, Gaithersburg, Maryland, U.S.A.*

<sup>‡</sup> *Department of Mathematics, Westminster College, 319 South Market Street, New  
Wilmington, Pennsylvania, U.S.A.*

---

## Abstract

In this paper we generalize and simplify generating functions for classical orthogonal polynomials, namely Jacobi, Gegenbauer, Chebyshev and Legendre polynomials. We generalize the generating function for Gegenbauer polynomials extended to Jacobi polynomials through examination of a two element sequence of generating functions for Jacobi polynomials. Simplifications of generating functions are accomplished through realization of general hypergeometric functions in terms of more elementary functions. We also use orthogonality relations for our classical orthogonal polynomials to determine definite integrals.

*Keywords:* Orthogonal polynomials, Generating functions, Gauss hypergeometric function, Eigenfunction expansions, Definite integrals  
*2000 MSC:* 33C45, 05A15, 33C05, 34L10, 30E20

---

## 1. Introduction

Orthogonal polynomials  $p_n$  (see Chapter 18 in ?) are polynomials which are orthogonal under an inner product on a real interval  $(a, b)$  with respect to some weight function  $w : (a, b) \rightarrow \mathbf{R}$ . In this paper we focus on so-called classical orthogonal polynomials. In our case we will focus on those orthogonal polynomials which are constructed by Gauss hypergeometric functions (Jacobi, Gegenbauer, Chebyshev, and Legendre polynomials). Since polynomials are entire functions on the complex plane, the study of orthogonal

polynomials extends itself naturally to complex analysis. Some fundamental texts which describe the properties of orthogonal polynomials include [1], [2], [3], and [4]. Our main focus in this paper is on generating functions for these classical orthogonal polynomials. Generating functions for classical orthogonal polynomials have a rich history. Many generating functions for the classical orthogonal polynomials can be found with tabulations and proofs in the above references on orthogonal polynomials. Some important references which focus particularly on the methods related to generating functions for orthogonal polynomials include [5] and [6].

A generating function  $F : (a, b) \times (0, 1) \rightarrow \mathbf{R}$  for an orthogonal polynomial is a series expansion

$$F(x, \rho) = \sum_{n=0}^{\infty} \alpha_n \rho^n p_n(x),$$

with coefficients  $\alpha_n$  independent of  $x$  and  $\rho$ . We refer to the (unique) generating function for an orthogonal polynomial when  $\alpha_n = 1$ . Generating functions for orthogonal polynomials can be used to define those orthogonal polynomials and are some of the most important functions known to be associated with those orthogonal polynomials.

In this paper, we use pre-existing generating functions for classical polynomials to simplify and/or generalize generating functions to construct alternative series expansions over the orthogonal polynomials which are no longer of generating function type. When possible we use the orthogonality property (see §18.2 in [7])

$$\int_a^b p_n(x) p_m(x) w(x) dx = h_n \delta_{n,m},$$

where  $\delta_{n,m} \in \{0, 1\}$  is the Kronecker delta, to obtain definite integral results from our alternative expansions. Unless stated otherwise the domains of convergence given in this paper are those of the original generating function and/or its corresponding definite integral.

Throughout this paper we rely on the following definitions. Let  $a_1, a_2, a_3, \dots \in \mathbf{C}$ . If  $i, j \in \mathbf{Z}$  and  $j < i$ , then  $\sum_{n=i}^j a_n = 0$  and  $\prod_{n=i}^j a_n = 1$ . The set of natural numbers is given by  $\mathbf{N} := \{1, 2, 3, \dots\}$ , the set  $\mathbf{N}_0 := \{0, 1, 2, \dots\} = \mathbf{N} \cup \{0\}$ , and  $\mathbf{Z} := \{0, \pm 1, \pm 2, \dots\}$ .

## 2. Jacobi polynomials

The Jacobi polynomials  $P_n^{(\alpha, \beta)} : \mathbf{C} \rightarrow \mathbf{C}$  can be defined in terms of the terminating Gauss hypergeometric series as follows ( (? , (18.5.7)))

$$P_n^{(\alpha, \beta)}(z) := \frac{(\alpha + 1)_n}{n!} {}_2F_1 \left( \begin{matrix} -n, n + \alpha + \beta + 1 \\ \alpha + 1 \end{matrix} ; \frac{1 - z}{2} \right),$$

for  $n \in \mathbf{N}_0$ , and  $\alpha, \beta > -1$ . The Gauss hypergeometric function  ${}_2F_1 : \mathbf{C}^2 \times (\mathbf{C} \setminus \mathbf{N}_0) \times \{z \in \mathbf{C} : |z| < 1\} \rightarrow \mathbf{C}$  (see Chapter 15 in ?) is defined as

$${}_2F_1 \left( \begin{matrix} a, b \\ c \end{matrix} ; z \right) := \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{z^n}{n!},$$

where the Pochhammer symbol (rising factorial)  $(\cdot)_n : \mathbf{C} \rightarrow \mathbf{C}$  (? , (5.2.4)) is defined by

$$(z)_n := \prod_{i=1}^n (z + i - 1),$$

where  $n \in \mathbf{N}_0$ . Note that the Gauss hypergeometric function can be analytically continued through for instance the Euler's integral representation for  $z \in \mathbf{C} \setminus (1, \infty)$  (see for instance (? , Theorem 2.2.1)). The orthogonality relation for Jacobi polynomials can be found in (? , (18.2.1), (18.2.5), Table 18.3.1)

$$\int_{-1}^1 P_m^{(\alpha, \beta)}(x) P_n^{(\alpha, \beta)}(x) (1-x)^\alpha (1+x)^\beta dx = \frac{2^{\alpha+\beta+1} \Gamma(\alpha + n + 1) \Gamma(\beta + n + 1)}{(2n + \alpha + \beta + 1) \Gamma(\alpha + \beta + n + 1) n!} \delta_{m,n}. \quad (1)$$

### 2.1. Expansions for Jacobi polynomials

Consider the generating function for Gegenbauer polynomials (see §3 for their definition) given by (? , (18.12.4)), namely

$$\frac{1}{(1 + \rho^2 - 2\rho x)^\nu} = \sum_{n=0}^{\infty} \rho^n C_n^\nu(x). \quad (2)$$

We attempt to generalize this expansion using the representation of Gegenbauer polynomials in terms of Jacobi polynomials given by (? , (18.7.1)), namely

$$C_n^\nu(x) = \frac{(2\nu)_n}{(\nu + \frac{1}{2})_n} P_n^{(\nu-1/2, \nu-1/2)}(x). \quad (3)$$

By making the replacement  $\nu - 1/2$  to either  $\alpha$  or  $\beta$ , we see that there are two possibilities for generalizing the Gegenbauer generating function to a generating function for Jacobi polynomials. The first is given in (?, (18.12.3))

$$\begin{aligned} & \frac{1}{(1+\rho)^{\alpha+\beta+1}} {}_2F_1 \left( \begin{matrix} \frac{\alpha+\beta+1}{2}, \frac{\alpha+\beta+2}{2} \\ \beta+1 \end{matrix} ; \frac{2\rho(1+x)}{(1+\rho)^2} \right) \\ &= \left( \frac{2}{\rho(1+x)} \right)^{\beta/2} \frac{\Gamma(\beta+1)}{(1+\rho^2-2\rho x)^{(\alpha+1)/2}} P_{\alpha}^{-\beta} \left( \frac{1+\rho}{\sqrt{1+\rho^2-2\rho x}} \right) \\ &= \sum_{n=0}^{\infty} \frac{(\alpha+\beta+1)_n}{(\beta+1)_n} \rho^n P_n^{(\alpha,\beta)}(x), \end{aligned} \quad (4)$$

where  $P_{\nu}^{\mu} : \mathbf{C} \setminus (-\infty, 1] \rightarrow \mathbf{C}$  is the associated Legendre function of the first kind (see Chapter 14 in ?), which can be defined in terms of the Gauss hypergeometric function as follows (?, (14.3.6), (15.2.2), §14.21(i))

$$P_{\nu}^{\mu}(z) := \frac{1}{\Gamma(1-\mu)} \left( \frac{z+1}{z-1} \right)^{\mu/2} {}_2F_1 \left( \begin{matrix} -\nu, \nu+1 \\ 1-\mu \end{matrix} ; \frac{1-z}{2} \right). \quad (5)$$

The associated Legendre function of the first kind can also be expressed in terms of the Gauss hypergeometric function as (see (?, (14.3.18), §14.21(iii))), namely

$$P_{\nu}^{\mu}(z) = \frac{2^{\mu} z^{\nu+\mu}}{\Gamma(1-\mu)(z^2-1)^{\mu/2}} {}_2F_1 \left( \begin{matrix} \frac{-\nu-\mu}{2}, \frac{-\nu-\mu+1}{2} \\ 1-\mu \end{matrix} ; 1 - \frac{1}{z^2} \right), \quad (6)$$

where  $|\arg(z-1)| < \pi$ . We have used (6) to simplify the generating function (4).

**Definition 1.** A **companion identity** is one which is produced by applying the map  $x \mapsto -x$  to an expansion over Jacobi polynomials or in terms of those orthogonal polynomials which can be obtained as limiting cases of Jacobi polynomials (i.e., Gegenbauer, Chebyshev, and Legendre polynomials) with argument  $x$  in conjunction with the parity relations for those orthogonal polynomials.

In our first generating function (4), we use the parity relation for Jacobi polynomials (see for instance (?, Table 18.6.1)), namely

$$P_n^{(\alpha,\beta)}(-x) = (-1)^n P_n^{(\beta,\alpha)}(x), \quad (7)$$

and the replacement  $\alpha, \beta \mapsto \beta, \alpha$  to produce the companion identity

$$\begin{aligned}
& \frac{1}{(1-\rho)^{\alpha+\beta+1}} {}_2F_1 \left( \begin{matrix} \frac{\alpha+\beta+1}{2}, \frac{\alpha+\beta+2}{2} \\ \alpha+1 \end{matrix} ; \frac{-2\rho(1-x)}{(1-\rho)^2} \right) \\
&= \left( \frac{2}{\rho(1-x)} \right)^{\alpha/2} \frac{\Gamma(\alpha+1)}{(1+\rho^2-2\rho x)^{(\beta+1)/2}} P_{\beta}^{-\alpha} \left( \frac{1-\rho}{\sqrt{1+\rho^2-2\rho x}} \right) \\
&= \sum_{n=0}^{\infty} \frac{(\alpha+\beta+1)_n}{(\alpha+1)_n} \rho^n P_n^{(\alpha,\beta)}(x), \quad (8)
\end{aligned}$$

where  $P_{\nu}^{\mu} : (-1, 1) \rightarrow \mathbf{C}$  is the Ferrers function of the first kind (associated Legendre function of the first kind on the cut) through (4.3.1), defined as

$$P_{\nu}^{\mu}(x) := \frac{1}{\Gamma(1-\mu)} \left( \frac{1+x}{1-x} \right)^{\mu/2} {}_2F_1 \left( \begin{matrix} -\nu, \nu+1 \\ 1-\mu \end{matrix} ; \frac{1-x}{2} \right). \quad (9)$$

The Ferrers function of the first kind can also be expressed in terms of the Gauss hypergeometric function as (see (4.3.2)), namely

$$P_{\nu}^{\mu}(x) = \frac{2^{\mu} x^{\nu+\mu}}{\Gamma(1-\mu)(1-x^2)^{\mu/2}} {}_2F_1 \left( \begin{matrix} \frac{-\nu-\mu}{2}, \frac{-\nu-\mu+1}{2} \\ 1-\mu \end{matrix} ; 1-\frac{1}{x^2} \right), \quad (10)$$

for  $x \in (0, 1)$ . We have used (10) to simplify (8).

There exists a natural extensions of (4), (8) (see (4.3.2)). The extension corresponding to (8) is given by

$$\begin{aligned}
& \frac{(\alpha+\beta+1)(1+\rho)}{(1-\rho)^{\alpha+\beta+2}} {}_2F_1 \left( \begin{matrix} \frac{\alpha+\beta+2}{2}, \frac{\alpha+\beta+3}{2} \\ \alpha+1 \end{matrix} ; \frac{-2\rho(1-x)}{(1-\rho)^2} \right) \\
&= \left( \frac{2}{\rho(1-x)} \right)^{\alpha/2} \frac{(\alpha+\beta+1)(1+\rho)\Gamma(\alpha+1)}{(1+\rho^2-2\rho x)^{(\beta+2)/2}} P_{\beta+1}^{-\alpha} \left( \frac{1-\rho}{\sqrt{1+\rho^2-2\rho x}} \right) \\
&= \sum_{n=0}^{\infty} (2n+\alpha+\beta+1) \frac{(\alpha+\beta+1)_n}{(\alpha+1)_n} \rho^n P_n^{(\alpha,\beta)}(x), \quad (11)
\end{aligned}$$

and its companion identity corresponding to (4) is

$$\begin{aligned}
& \frac{(\alpha + \beta + 1)(1 - \rho)}{(1 + \rho)^{\alpha + \beta + 2}} {}_2F_1 \left( \begin{matrix} \frac{\alpha + \beta + 2}{2}, \frac{\alpha + \beta + 3}{2} \\ \beta + 1 \end{matrix}; \frac{2\rho(1 + x)}{(1 + \rho)^2} \right) \\
&= \left( \frac{2}{\rho(1 + x)} \right)^{\beta/2} \frac{(\alpha + \beta + 1)(1 - \rho)\Gamma(\beta + 1)}{(1 + \rho^2 - 2\rho x)^{(\alpha + 2)/2}} P_{\alpha + 1}^{-\beta} \left( \frac{1 + \rho}{\sqrt{1 + \rho^2 - 2\rho x}} \right) \\
&= \sum_{n=0}^{\infty} (2n + \alpha + \beta + 1) \frac{(\alpha + \beta + 1)_n}{(\beta + 1)_n} \rho^n P_n^{(\alpha, \beta)}(x). \quad (12)
\end{aligned}$$

We have used (6), (10) to simplify these Gauss hypergeometric function generating functions as associated Legendre functions.

Upon examination of these two sets of generating functions, we suspected that these were just two examples of an infinite sequence of such expansions. This led us to the proof of the following theorem, which is a Jacobi polynomial expansion which generalizes the generating function for Gegenbauer polynomials (2).

**Theorem 1.** *Let  $m \in \mathbf{N}_0$ ,  $\alpha, \beta > -1$ ,  $x \in [-1, 1]$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\} \setminus (-1, 0]$ . Then*

$$\begin{aligned}
& \frac{(1 + x)^{-\beta/2}}{(1 + \rho^2 - 2\rho x)^{(\alpha + m + 1)/2}} P_{\alpha + m}^{-\beta} \left( \frac{1 + \rho}{\sqrt{1 + \rho^2 - 2\rho x}} \right) \\
&= \frac{\rho^{-(\alpha + 1)/2}}{2^{\beta/2}(1 - \rho)^m} \sum_{n=0}^{\infty} \frac{(2n + \alpha + \beta + 1)\Gamma(\alpha + \beta + n + 1)(\alpha + \beta + m + 1)_{2n}}{\Gamma(\beta + n + 1)} \\
&\quad \times P_{-m}^{-\alpha - \beta - 2n - 1} \left( \frac{1 + \rho}{1 - \rho} \right) P_n^{(\alpha, \beta)}(x). \quad (13)
\end{aligned}$$

**Proof.** Let  $\rho \in (0, \epsilon)$  with  $\epsilon$  sufficiently small. Then using the definition of the following Gauss hypergeometric function

$$\begin{aligned}
& {}_2F_1 \left( \begin{matrix} \frac{\alpha + \beta + m + 1}{2}, \frac{\alpha + \beta + m + 2}{2} \\ \alpha + 1 \end{matrix}; \frac{-2\rho(1 - x)}{(1 - \rho)^2} \right) \\
&= \sum_{n=0}^{\infty} \frac{\left(\frac{\alpha + \beta + m + 1}{2}\right)_n \left(\frac{\alpha + \beta + m + 2}{2}\right)_n (-2\rho)^n (1 - x)^n}{n!(\alpha + 1)_n (1 - \rho)^{2n}}, \quad (14)
\end{aligned}$$

the expansion of  $(1-x)^n$  in terms of Jacobi polynomials is given by

$$(1-x)^n = 2^n(\alpha+1)_n \sum_{k=0}^n \frac{(-n)_k(\alpha+\beta+2k+1)(\alpha+\beta+1)_k}{(\alpha+\beta+1)_{n+k+1}(\alpha+1)_k} P_k^{(\alpha,\beta)}(x), \quad (15)$$

whose coefficients can be determined using orthogonality of Jacobi polynomials (1) combined with the Mellin transform given in (?, (18.17.36)). By inserting (15) in the right-hand side of (14), we obtain an expansion of the Gauss hypergeometric function on the left-hand side in terms of Jacobi polynomials. By interchanging the two sums (with justification by absolute convergence) we are left with a Gauss hypergeometric function as a coefficient of the Jacobi polynomial expansion. The resulting expansion formula is

$$\begin{aligned} & \frac{(\alpha+\beta+1)_m(1-\rho)^m}{(1-\rho)^{\alpha+\beta+m+1}} {}_2F_1 \left( \begin{matrix} \frac{\alpha+\beta+m+1}{2}, \frac{\alpha+\beta+m+2}{2} \\ \beta+1 \end{matrix}; \frac{2\rho(1+x)}{(1+\rho)^2} \right) \\ = & \sum_{n=0}^{\infty} (2n+\alpha+\beta+1)_m \frac{(\alpha+\beta+1)_n}{(\beta+1)_n} {}_2F_1 \left( \begin{matrix} m, -m+1 \\ \alpha+\beta+2n+2 \end{matrix}; \frac{-\rho}{1-\rho} \right) \rho^n P_n^{(\alpha,\beta)}(x). \end{aligned} \quad (16)$$

The Gauss hypergeometric function coefficient of the expansion is realized to be an associated Legendre function of the first kind through (5). The Gauss hypergeometric function on the left-hand of (16) can be simplified as well using (6). Now we apply the expansion theorem for Jacobi polynomials (see for instance (?, Theorem 9.1.1, (9.2.3))). The Jacobi polynomial expansion is equivalent to the integral for the coefficients and both sides of (13) are analytic functions for  $\rho \in \{z \in \mathbf{C} : |z| < 1\} \setminus (-1, 0]$ . Since we know that (13) is valid for  $\rho \in (0, \epsilon)$ , then by the identity theorem for analytic functions, the equation hold for this region. This completes the proof.  $\blacksquare$

We have also derived the companion identity to (13), which we give in the following corollary.

**Corollary 1.** *Let  $m \in \mathbf{N}_0$ ,  $\alpha, \beta > -1$ ,  $x \in [-1, 1]$ ,  $\rho \in (0, 1)$ . Then*

$$\begin{aligned} & \frac{(1-x)^{-\alpha/2}}{(1+\rho^2-2\rho x)^{(\beta+m+1)/2}} P_{\beta+m}^{-\alpha} \left( \frac{1-\rho}{\sqrt{1+\rho^2-2\rho x}} \right) \\ &= \frac{\rho^{-(\beta+1)/2}}{2^{\alpha/2}(1+\rho)^m} \sum_{n=0}^{\infty} \frac{(2n+\alpha+\beta+1)\Gamma(\alpha+\beta+n+1)(\alpha+\beta+m+1)_{2n}}{\Gamma(\alpha+n+1)} \\ & \quad \times P_{-m}^{-\alpha-\beta-2n-1} \left( \frac{1-\rho}{1+\rho} \right) P_n^{(\alpha,\beta)}(x). \end{aligned} \quad (17)$$

**Proof.** We start with (16) and apply the parity relation for Jacobi polynomials (7). Let  $\rho \in (0, 1)$ . The Gauss hypergeometric function coefficient of the Jacobi expansion is seen to be a Ferrers function of the first kind (9). After the application of the parity relation, the left-hand side also reduces to a Ferrers function of the first kind through (10). This completes the proof. ■

Theorem 1 generalizes (4), (12), while Corollary 1 generalizes (8), (11). Both Theorem 1 and Corollary 1 generalize the generating function for Gegenbauer polynomials (2) since it is easily verified to be its own companion identity. According to ?, (11), (12), are closely related to the Poisson kernel for Jacobi polynomials, so we hope our generalizations have applications as such.

We then looked for other generating functions of Jacobi polynomials that could be simplified. The following two theorems originate from ?, and are for generating functions of Jacobi polynomials with even and odd degree respectively. In order to prove the next two theorems, it will be necessary to introduce (and prove) the following two lemmas.

**Lemma 1.** *Let  $\alpha, w \in \mathbf{C}$ . Then*

$${}_2F_1\left(\begin{matrix} \frac{-\alpha}{2}, \frac{-\alpha+1}{2} \\ \frac{1}{2} \end{matrix}; w^2\right) = \frac{1}{2} [(1+w)^\alpha + (1-w)^\alpha].$$

**Proof.** Using entry 11 in (?, p. 157) for the associated Legendre function of the first kind (6) along with (?, (8.6.8)), namely

$$P_\nu^{1/2}(z) = \frac{1}{\sqrt{2\pi}} (z^2 - 1)^{-1/4} \left[ \left( z + \sqrt{z^2 - 1} \right)^{\nu+1/2} + \left( z - \sqrt{z^2 - 1} \right)^{-\nu-1/2} \right], \quad (18)$$

and (?, (14.5.11)), to simplify the Gauss hypergeometric function. Note that if  $w$  lies along the imaginary axis then one should use the corresponding Ferrers function representations (10). ■

**Lemma 2.** *Let  $\alpha, w \in \mathbf{C} \setminus \{0\}$ . Then*

$${}_2F_1\left(\begin{matrix} \frac{-\alpha+1}{2}, \frac{-\alpha+2}{2} \\ \frac{3}{2} \end{matrix}; w^2\right) = \frac{1}{2\alpha w} [(1+w)^\alpha - (1-w)^\alpha].$$

**Proof.** Using (6), along with (?, (8.6.9)), namely

$$P_\nu^{-1/2}(z) = \sqrt{\frac{2}{\pi}} \frac{(z^2 - 1)^{-1/4}}{(2\nu + 1)} \left[ \left( z + \sqrt{z^2 - 1} \right)^{\nu+1/2} - \left( z + \sqrt{z^2 - 1} \right)^{-\nu-1/2} \right], \quad (19)$$

and (?, (14.5.12)), to simplify the Gauss hypergeometric function. Note once more that if  $w$  lies along the imaginary axis then one should use the corresponding Ferrers function representations. This completes the proof. ■

**Theorem 2.** Let  $\alpha, \beta > -1$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in [-1, 1]$ . Then

$$\begin{aligned} \left( 1 + \frac{(1+x)\sqrt{\rho}}{2} \right)^\alpha \left( 1 - \frac{(1-x)\sqrt{\rho}}{2} \right)^\beta + \left( 1 - \frac{(1+x)\sqrt{\rho}}{2} \right)^\alpha \left( 1 + \frac{(1-x)\sqrt{\rho}}{2} \right)^\beta \\ = 2 \sum_{n=0}^{\infty} \rho^n P_{2n}^{(\alpha-2n, \beta-2n)}(x). \end{aligned} \quad (20)$$

**Proof.** In problem 11, p. 215 of ? there is a generating function for Jacobi polynomials, namely

$$\begin{aligned} {}_2F_1 \left( \begin{matrix} -\frac{\alpha}{2}, \frac{-\alpha+1}{2} \\ \frac{1}{2} \end{matrix}; \frac{(1+x)^2 \rho}{4} \right) {}_2F_1 \left( \begin{matrix} -\frac{\beta}{2}, \frac{-\beta+1}{2} \\ \frac{1}{2} \end{matrix}; \frac{(1-x)^2 \rho}{4} \right) \\ + \frac{1}{4} \alpha \beta (x^2 - 1) \rho {}_2F_1 \left( \begin{matrix} \frac{-\alpha+1}{2}, \frac{-\alpha+2}{2} \\ \frac{3}{2} \end{matrix}; \frac{(1+x)^2 \rho}{4} \right) {}_2F_1 \left( \begin{matrix} \frac{-\beta+1}{2}, \frac{-\beta+2}{2} \\ \frac{3}{2} \end{matrix}; \frac{(1-x)^2 \rho}{4} \right) \\ = \sum_{n=0}^{\infty} \rho^n P_{2n}^{(\alpha-2n, \beta-2n)}(x). \end{aligned} \quad (21)$$

Applying Lemmas 1 and 2 with either  $\nu = \alpha - 1/2, w = (1+x)\sqrt{\rho}/2$  or  $\nu = \beta - 1/2, w = (1-x)\sqrt{\rho}/2$  for the Gauss hypergeometric functions on the left-hand side of (21) and simplifying completes the proof. ■

**Theorem 3.** Let  $\alpha, \beta > -1$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in [-1, 1]$ . Then

$$\begin{aligned} \left( 1 + \frac{(1+x)\sqrt{\rho}}{2} \right)^{\alpha+1} \left( 1 - \frac{(1-x)\sqrt{\rho}}{2} \right)^{\beta+1} - \left( 1 - \frac{(1+x)\sqrt{\rho}}{2} \right)^{\alpha+1} \left( 1 + \frac{(1-x)\sqrt{\rho}}{2} \right)^{\beta+1} \\ = 2\sqrt{\rho} \sum_{n=0}^{\infty} \rho^n P_{2n+1}^{(\alpha-2n, \beta-2n)}(x). \end{aligned} \quad (22)$$

**Proof.** In problem 11, p. 215 of ? there is a generating function for Jacobi polynomials (Note that the generating function is stated wrong in ?, the last  ${}_2F_1$  should have a  $3/2$  instead of a  $1/2$ ), namely

$$\begin{aligned} & \frac{(\alpha+1)(x+1)}{2} {}_2F_1 \left( \begin{matrix} -\frac{\alpha}{2}, \frac{-\alpha+1}{2} \\ \frac{3}{2} \end{matrix}; \frac{(1+x)^2\rho}{4} \right) {}_2F_1 \left( \begin{matrix} \frac{-\beta-1}{2}, -\frac{\beta}{2} \\ \frac{1}{2} \end{matrix}; \frac{(1-x)^2\rho}{4} \right) \\ & + \frac{(\beta+1)(x-1)}{2} {}_2F_1 \left( \begin{matrix} \frac{-\alpha-1}{2}, -\frac{\alpha}{2} \\ \frac{1}{2} \end{matrix}; \frac{(1+x)^2\rho}{4} \right) {}_2F_1 \left( \begin{matrix} -\frac{\beta}{2}, \frac{-\beta+1}{2} \\ \frac{3}{2} \end{matrix}; \frac{(1-x)^2\rho}{4} \right) \\ & = \sum_{n=0}^{\infty} \rho^n P_{2n+1}^{(\alpha-2n, \beta-2n)}(x). \end{aligned} \quad (23)$$

Applying Lemmas 1 and 2 with either  $\nu = \alpha + 1/2, w = (1+x)\sqrt{\rho}/2$  or  $\nu = \beta + 1/2, w = (1-x)\sqrt{\rho}/2$  for the Gauss hypergeometric functions on the left-hand side of (23) and simplifying completes the proof.  $\blacksquare$

It is straightforward to show that the parity relation for Jacobi polynomials (7) here creates companion identities that are the same as the two previous derived theorems.

Furthermore, it should be stated that Theorems 2, 3, as well as all the corollaries associated with these theorems below, are consistent with (?, cf. (8)), namely

$$\left(1 + \frac{(1+x)\rho}{2}\right)^{\alpha} \left(1 - \frac{(1-x)\rho}{2}\right)^{\beta} = \sum_{n=0}^{\infty} \rho^n P_n^{(\alpha-n, \beta-n)}(x),$$

through the parity relation for Jacobi polynomials (7).

## 2.2. Definite integrals for Jacobi polynomials

**Corollary 2.** Let  $m, n \in \mathbf{N}_0$ ,  $\alpha, \beta > -1$ ,  $\rho \in \{z \in \mathbf{C} : 0 < |z| < 1\} \setminus (-1, 0]$ .

Then

$$\begin{aligned} & \int_{-1}^1 \frac{(1-x)^{\alpha}(1+x)^{\beta/2}}{(1+\rho^2-2\rho x)^{(\alpha+m+1)/2}} P_{\alpha+m}^{-\beta} \left( \frac{1+\rho}{\sqrt{1+\rho^2-2\rho x}} \right) P_n^{(\alpha, \beta)}(x) dx \\ & = \frac{2^{\alpha+\beta/2+1} \Gamma(\alpha+n+1)(\alpha+\beta+m+1)_{2n}}{n! \rho^{(\alpha+1)/2} (1-\rho)^m} P_{-m}^{-\alpha-\beta-2n-1} \left( \frac{1+\rho}{1-\rho} \right). \end{aligned}$$

**Proof.** Multiply both sides of (13) by  $(1-x)^\alpha(1+x)^\beta P_k^{(\alpha,\beta)}(x)$ , for  $k \in \mathbf{N}_0$  and integrate over  $(-1, 1)$  using the orthogonality relation (1). ■

**Corollary 3.** *Let  $m, n \in \mathbf{N}_0$ ,  $\alpha, \beta > -1$ ,  $\rho \in (0, 1)$ . Then*

$$\begin{aligned} \int_{-1}^1 \frac{(1-x)^{\alpha/2}(1+x)^\beta}{(1+\rho^2-2\rho x)^{(\beta+m+1)/2}} P_{\beta+m}^{-\alpha} \left( \frac{1-\rho}{\sqrt{1+\rho^2-2\rho x}} \right) P_n^{(\alpha,\beta)}(x) dx \\ = \frac{2^{\alpha/2+\beta+1} \Gamma(\beta+n+1)(\alpha+\beta+m+1)_{2n}}{n! \rho^{(\beta+1)/2} (1+\rho)^m} P_{-m}^{-\alpha-\beta-2n-1} \left( \frac{1-\rho}{1+\rho} \right). \end{aligned}$$

**Proof.** Same as in the proof of Corollary 2, except apply to both sides of (17). ■

### 2.3. Expansions and definite integrals from the Szegő transformation

If one applies on the complex plane, the Szegő transformation (conformal map)

$$z = \frac{1+\rho^2}{2\rho}, \quad (24)$$

(which maps a circle with radius less than unity to an ellipse with foci at  $\pm 1$ ) to the expansion in Theorem 1, then one obtains a new expansion. By (? , Theorem 12.7.3), this new Jacobi polynomial expansion is convergent for all  $x \in \mathbf{C}$  within the interior of this ellipse. Applying (24) to (13) yields the following corollary.

**Corollary 4.** *Let  $m \in \mathbf{N}_0$ ,  $\alpha, \beta > -1$ ,  $x, z \in \mathbf{C}$ , with  $z \in \mathbf{C} \setminus (-\infty, 1]$  on any ellipse with foci at  $\pm 1$  and  $x$  in the interior of that ellipse. Then*

$$\begin{aligned} \frac{(1+x)^{-\beta/2}}{(z-x)^{(\alpha+m+1)/2}} P_{\alpha+m}^{-\beta} \left( \frac{1+z-\sqrt{z^2-1}}{\sqrt{2(z-\sqrt{z^2-1})(z-x)}} \right) \\ = 2^{(\alpha-\beta+m+1)/2} \frac{(z-\sqrt{z^2-1})^{m/2}}{(1-z+\sqrt{z^2-1})^m} \sum_{n=0}^{\infty} \frac{(2n+\alpha+\beta+1)\Gamma(\alpha+\beta+n+1)(\alpha+\beta+m+1)_{2n}}{\Gamma(\beta+n+1)} \\ \times P_{-m}^{-\alpha-\beta-2n-1} \left( \sqrt{\frac{z+1}{z-1}} \right) P_n^{(\alpha,\beta)}(x). \end{aligned} \quad (25)$$

We would just like to briefly note that one may use the Szegő transformation (24) to obtain new expansion formulae and corresponding definite integrals from all the Jacobi, Gegenbauer, Legendre and Chebyshev polynomial expansions used in this paper. For the sake of brevity, we leave this to the reader.

### 3. Gegenbauer polynomials

The Gegenbauer polynomials  $C_n^\mu : \mathbf{C} \rightarrow \mathbf{C}$  can be defined in terms of the terminating Gauss hypergeometric series as follows ( (? , (18.5.9)))

$$C_n^\mu(z) := \frac{(2\mu)_n}{n!} {}_2F_1 \left( \begin{matrix} -n, n+2\mu \\ \mu + \frac{1}{2} \end{matrix} ; \frac{1-z}{2} \right), \quad (26)$$

for  $n \in \mathbf{N}_0$  and  $\mu \in (-1/2, \infty) \setminus \{0\}$ . The orthogonality relation for Gegenbauer polynomials can be found in (? , (18.2.1), (18.2.5), Table 18.3.1) for  $m, n \in \mathbf{N}_0$  and  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,

$$\int_{-1}^1 C_m^\mu(x) C_n^\mu(x) (1-x^2)^{\mu-1/2} dx = \frac{\pi 2^{1-2\mu} \Gamma(2\mu+n)}{(n+\mu) \Gamma^2(\mu) n!} \delta_{m,n}. \quad (27)$$

The Gegenbauer polynomials (26) are defined for  $\mu \in (-1/2, \infty) \setminus \{0\}$ . However many of the formulae listed below actually make sense in the limit as  $\mu \rightarrow 0$ . In this case, one should take the limit of the expression as  $\mu \rightarrow 0$  with the interpretation of obtaining Chebyshev polynomials of the first kind (see §6 for the details of this limiting procedure).

#### 3.1. Expansions for Gegenbauer polynomials

**Corollary 5.** *Let  $m \in \mathbf{N}_0$ ,  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\} \setminus (-1, 0]$ ,  $x \in [-1, 1]$ . Then*

$$\begin{aligned} & \frac{1}{(1+\rho^2-2\rho x)^{\mu+m/2}} C_m^\mu \left( \frac{1+\rho}{\sqrt{1+\rho^2-2\rho x}} \right) \\ &= \frac{2\Gamma(2\mu+m)}{m! \rho^\mu (1-\rho)^m} \sum_{n=0}^{\infty} (n+\mu)(2\mu+m) {}_2F_{-m}^{-2\mu-2n} \left( \frac{1+\rho}{1-\rho} \right) C_n^\mu(x). \end{aligned} \quad (28)$$

**Proof.** Using (13), substituting  $\alpha = \beta = \mu - 1/2$  along with (3) and (? , (14.3.22)), namely

$$P_{n+\mu-1/2}^{1/2-\mu}(z) = \frac{2^{\mu-1/2}\Gamma(\mu)n!}{\sqrt{\pi}\Gamma(2\mu+n)}(z^2-1)^{\mu/2-1/4}C_n^\mu(z),$$

completes the proof. ■

**Corollary 6.** Let  $m \in \mathbf{N}_0$ ,  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in (0, 1)$ ,  $x \in [-1, 1]$ . Then

$$\begin{aligned} & \frac{1}{(1+\rho^2-2\rho x)^{\mu+m/2}}C_m^\mu\left(\frac{1-\rho}{\sqrt{1+\rho^2-2\rho x}}\right) \\ &= \frac{2\Gamma(2\mu+m)}{m!\rho^\mu(1+\rho)^m}\sum_{n=0}^{\infty}(n+\mu)(2\mu+m)_{2n}P_{-m}^{-2\mu-2n}\left(\frac{1-\rho}{1+\rho}\right)C_n^\mu(x). \end{aligned} \quad (29)$$

**Proof.** Using (17), substituting  $\alpha = \beta = \mu - 1/2$  along with (3) and (? , (14.3.21)), namely

$$P_{n+\mu-1/2}^{1/2-\mu}(x) = \frac{2^{\mu-1/2}\Gamma(\mu)n!}{\sqrt{\pi}\Gamma(2\mu+n)}(1-x^2)^{\mu/2-1/4}C_n^\mu(x),$$

completes the proof. ■

**Theorem 4.** Let  $\alpha > -1$ ,  $\alpha \notin \{3/2, 5/2, 7/2, \dots\}$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in [-1, 1]$ . Then

$$\begin{aligned} & \left(1 + \sqrt{\rho}x - \frac{(1-x^2)\rho}{4}\right)^\alpha + \left(1 - \sqrt{\rho}x - \frac{(1-x^2)\rho}{4}\right)^\alpha \\ &= 2 \sum_{n=0}^{\infty} \frac{(-\alpha + \frac{1}{2})_n (-\alpha)_n}{2^{4n} \left(\frac{-\alpha}{2} + \frac{3}{4}\right)_n \left(\frac{-\alpha}{2} + \frac{1}{4}\right)_n} \rho^n C_{2n}^{\alpha+1/2-2n}(x). \end{aligned}$$

When  $\alpha = 1/2$ , then

$$\begin{aligned} & \sqrt{1 + \sqrt{\rho}x - \frac{(1-x^2)\rho}{4}} + \sqrt{1 - \sqrt{\rho}x - \frac{(1-x^2)\rho}{4}} \\ &= - \sum_{n=0}^{\infty} \frac{2^{2-4n}}{(2n-1)(1+\delta_{n,0})} \rho^n C_{2n}^{1-2n}(x). \end{aligned}$$

**Proof.** Using (3) in (20) with simplification completes the proof. The restriction on  $\alpha$  comes from the Pochhammer symbols in the denominator. When  $\alpha \in \{3/2, 5/2, \dots\}$ , one of the Pochhammer symbols is zero at some value of  $n$ , and the sum becomes indeterminate. When  $\alpha = 1/2$ , it is possible to express the Pochhammer symbols as above to complete the proof. ■

**Theorem 5.** Let  $\alpha > -1$ ,  $\alpha \notin \{3/2, 5/2, 7/2, \dots\}$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in [-1, 1]$ . Then

$$\begin{aligned} & \left(1 + \sqrt{\rho}x - \frac{(1-x^2)\rho}{4}\right)^{\alpha+1} - \left(1 - \sqrt{\rho}x - \frac{(1-x^2)\rho}{4}\right)^{\alpha+1} \\ &= \frac{(2\alpha+2)\sqrt{\rho}}{2\alpha+1} \sum_{n=0}^{\infty} \frac{(-\alpha)_n (-\alpha - \frac{1}{2})_n}{2^{4n} \left(\frac{-\alpha}{2} + \frac{3}{4}\right)_n \left(\frac{-\alpha}{2} + \frac{1}{4}\right)_n} \rho^n C_{2n+1}^{\alpha+1/2-2n}(x). \end{aligned}$$

**Proof.** Same as proof of Theorem 4 except apply to (22). ■

**Theorem 6.** Let  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in \{z \in \mathbf{C} : z \text{ interior to an ellipse with foci at } \pm 1 \text{ with semi-major axis } (1+\rho)/(2\sqrt{\rho})\}$ . Then

$$\frac{1}{(1+\rho-2\sqrt{\rho}x)^\mu} + \frac{1}{(1+\rho+2\sqrt{\rho}x)^\mu} = 2 \sum_{n=0}^{\infty} \rho^n C_{2n}^\mu(x). \quad (30)$$

**Proof.** In problem 5, p. 212 of ? there is a generating function for Gegenbauer polynomials, namely

$$\begin{aligned} & {}_2F_1\left(\begin{matrix} \frac{\mu}{2}, \frac{\mu+1}{2} \\ \frac{1}{2} \end{matrix}; \rho \left(x - \sqrt{x^2-1}\right)^2\right) {}_2F_1\left(\begin{matrix} \frac{\mu}{2}, \frac{\mu+1}{2} \\ \frac{1}{2} \end{matrix}; \rho \left(x + \sqrt{x^2-1}\right)^2\right) \\ & + \mu^2 \rho {}_2F_1\left(\begin{matrix} \frac{\mu+1}{2}, \frac{\mu+2}{2} \\ \frac{3}{2} \end{matrix}; \rho \left(x - \sqrt{x^2-1}\right)^2\right) {}_2F_1\left(\begin{matrix} \frac{\mu+1}{2}, \frac{\mu+2}{2} \\ \frac{3}{2} \end{matrix}; \rho \left(x + \sqrt{x^2-1}\right)^2\right) \\ & = \sum_{n=0}^{\infty} \rho^n C_{2n}^\mu(x). \end{aligned} \quad (31)$$

Using Lemmas 1 and 2 with  $\nu = -\mu - 1/2$  and either  $w = \sqrt{\rho}(x - \sqrt{x^2-1})$  or  $w = \sqrt{\rho}(x + \sqrt{x^2-1})$  for the Gauss hypergeometric functions on the

left-hand side of (31) and simplifying produces the result. The domain of convergence for  $x$  follows from the fact that  $|x \pm \sqrt{x^2 - 1}| = \sigma > 1$  is an ellipse with foci at  $\pm 1$  in the complex plane with semi-major axis given by  $(\sigma^2 + 1)/(2\sigma)$ , and for this domain the arguments of the Gauss hypergeometric functions are less than unity. ■

**Theorem 7.** *Let  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in \{z \in \mathbf{C} : z \text{ interior to an ellipse with foci at } \pm 1 \text{ with semi-major axis } (1 + \rho)/(2\sqrt{\rho})\}$ . Then*

$$\frac{1}{(1 + \rho - 2\sqrt{\rho}x)^\mu} - \frac{1}{(1 + \rho + 2\sqrt{\rho}x)^\mu} = 2\sqrt{\rho} \sum_{n=0}^{\infty} \rho^n C_{2n+1}^\mu(x). \quad (32)$$

**Proof.** In problem 5, p. 213 of ? there is a generating function for Gegenbauer polynomials, namely

$$\begin{aligned} & \frac{\mu}{x + \sqrt{x^2 - 1}} {}_2F_1 \left( \begin{matrix} \frac{\mu}{2}, \frac{\mu+1}{2} \\ \frac{1}{2} \end{matrix} ; \rho \left( x + \sqrt{x^2 - 1} \right)^2 \right) {}_2F_1 \left( \begin{matrix} \frac{\mu+1}{2}, \frac{\mu+2}{2} \\ \frac{3}{2} \end{matrix} ; \rho \left( x - \sqrt{x^2 - 1} \right)^2 \right) \\ & + \frac{\mu}{x - \sqrt{x^2 - 1}} {}_2F_1 \left( \begin{matrix} \frac{\mu}{2}, \frac{\mu+1}{2} \\ \frac{1}{2} \end{matrix} ; \rho \left( x - \sqrt{x^2 - 1} \right)^2 \right) {}_2F_1 \left( \begin{matrix} \frac{\mu+1}{2}, \frac{\mu+2}{2} \\ \frac{3}{2} \end{matrix} ; \rho \left( x + \sqrt{x^2 - 1} \right)^2 \right) \\ & = \sum_{n=0}^{\infty} \rho^n C_{2n+1}^\mu(x). \quad (33) \end{aligned}$$

Using Lemmas 1 and 2 with  $\nu = -\mu - 1/2$  and either  $w = \sqrt{\rho}/(x + \sqrt{x^2 - 1})$  or  $w = \sqrt{\rho}/(x - \sqrt{x^2 - 1})$  for the Gauss hypergeometric functions on the left-hand side of (33) and simplifying completes the proof. The proof of Theorem 6 contains a corresponding explanation for the domain of convergence for  $x$ . ■

It should be stated that Theorems 6, 7, as well as all the corollaries associated with these theorems below, are consistent with the generating function for Gegenbauer polynomials (2) through the parity relation for Gegenbauer polynomials, cf. (7) and (?, (18.7.1)).

**Theorem 8.** Let  $\lambda \in \mathbf{C}$ ,  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in (0, 1)$ ,  $x \in [-1, 1]$ . Then

$$\begin{aligned} (1-x^2)^{1/4-\mu/2} P_{\mu-\lambda-1/2}^{1/2-\mu} \left( \sqrt{1+\rho^2-2\rho x} + \rho \right) P_{\mu-\lambda-1/2}^{1/2-\mu} \left( \sqrt{1+\rho^2-2\rho x} - \rho \right) \\ = \frac{2^{1/2-\mu}}{\Gamma(\mu + \frac{1}{2})} \sum_{n=0}^{\infty} \frac{(\lambda)_n (2\mu - \lambda)_n}{(2\mu)_n \Gamma(\mu + \frac{1}{2} + n)} \rho^{\mu-1/2+n} C_n^\mu(x). \end{aligned} \quad (34)$$

**Proof.** (?, (9.8.32)) give a generating function for Gegenbauer polynomials, namely

$$\begin{aligned} {}_2F_1 \left( \begin{matrix} \lambda, 2\mu - \lambda \\ \mu + \frac{1}{2} \end{matrix}; \frac{1 - \sqrt{1+\rho^2-2\rho x} - \rho}{2} \right) {}_2F_1 \left( \begin{matrix} \lambda, 2\mu - \lambda \\ \mu + \frac{1}{2} \end{matrix}; \frac{1 - \sqrt{1+\rho^2-2\rho x} + \rho}{2} \right) \\ = \sum_{n=0}^{\infty} \frac{(\lambda)_n (2\mu - \lambda)_n}{(2\mu)_n (\mu + \frac{1}{2})_n} \rho^n C_n^\mu(x). \end{aligned}$$

Using (?, (15.8.17)) to do a quadratic transformation on the Gauss hypergeometric functions and then using (6), with degree and order given by  $\mu - \lambda - 1/2$ ,  $1/2 - \mu$ , respectively and either  $z = \sqrt{1+\rho^2-2\rho x} + \rho$  or  $z = \sqrt{1+\rho^2-2\rho x} - \rho$ . Simplification completes the proof. ■

**Theorem 9.** Let  $\alpha \in \mathbf{C}$ ,  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in (0, 1)$ ,  $x \in [-1, 1]$ . Then

$$\begin{aligned} \frac{(1-x^2)^{1/4-\mu/2}}{(1+\rho^2-2\rho x)^{1/4+\alpha/2-\mu/2}} P_{\mu-\alpha-1/2}^{1/2-\mu} \left( \frac{1-\rho x}{\sqrt{1+\rho^2-2\rho x}} \right) \\ = \frac{(\rho/2)^{\mu-1/2}}{\Gamma(\mu + \frac{1}{2})} \sum_{n=0}^{\infty} \frac{(\alpha)_n}{(2\mu)_n} \rho^n C_n^\mu(x). \end{aligned} \quad (35)$$

**Proof.** On p. 279 of (?, (8)) there is a generating function for Gegenbauer polynomials, namely

$$(1-\rho x)^{-\alpha} {}_2F_1 \left( \begin{matrix} \frac{\alpha}{2}, \frac{\alpha+1}{2} \\ \mu + \frac{1}{2} \end{matrix}; \frac{-\rho^2(1-x^2)}{(1-\rho x)^2} \right) = \sum_{n=0}^{\infty} \frac{(\alpha)_n}{(2\mu)_n} \rho^n C_n^\mu(x).$$

Using (6) to rewrite the Gauss hypergeometric function on the left-hand side of the above equation completes the proof. ■

### 3.2. Definite integrals for Gegenbauer polynomials

**Corollary 7.** Let  $m, n \in \mathbf{N}_0$ ,  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in \{z \in \mathbf{C} : 0 < |z| < 1\} \setminus (-1, 0]$ . Then

$$\begin{aligned} \int_{-1}^1 \frac{(1-x^2)^{\mu-1/2}}{(1+\rho^2-2\rho x)^{m/2+\mu}} C_m^\mu \left( \frac{1+\rho}{\sqrt{1+\rho^2-2\rho x}} \right) C_n^\mu(x) dx \\ = \frac{2^{2-2\mu} \pi \Gamma(2\mu+n) \Gamma(2\mu+2n+m)}{m!n! \Gamma^2(\mu) \rho^\mu (1-\rho)^m} P_{-m}^{-2n-2\mu} \left( \frac{1+\rho}{1-\rho} \right). \end{aligned}$$

**Proof.** Multiply both sides of (28) by  $(1-x^2)^{\mu-1/2} C_k^\mu(x)$ , for  $k \in \mathbf{N}_0$  and integrate over  $(-1, 1)$  using the orthogonality relation (27). ■

**Corollary 8.** Let  $m, n \in \mathbf{N}_0$ ,  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in (0, 1)$ . Then

$$\begin{aligned} \int_{-1}^1 \frac{(1-x^2)^{\mu-1/2}}{(1+\rho^2-2\rho x)^{m/2+\mu}} C_m^\mu \left( \frac{1-\rho}{\sqrt{1+\rho^2-2\rho x}} \right) C_n^\mu(x) dx \\ = \frac{2^{2-2\mu} \pi \Gamma(2\mu+n) \Gamma(2\mu+2n+m)}{m!n! \Gamma^2(\mu) \rho^\mu (1+\rho)^m} P_{-m}^{-2n-2\mu} \left( \frac{1-\rho}{1+\rho} \right). \end{aligned}$$

**Proof.** Same as in the proof of Corollary 7, except apply to both sides of (29). ■

**Corollary 9.** Let  $n \in \mathbf{N}_0$ ,  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ . Then

$$\begin{aligned} \int_{-1}^1 (1-x^2)^{\mu-1/2} \left( \frac{1}{(1+\rho-2\sqrt{\rho}x)^\mu} + \frac{1}{(1+\rho+2\sqrt{\rho}x)^\mu} \right) C_n^\mu(x) dx \\ = \begin{cases} \frac{2\sqrt{\pi} \Gamma(\mu+\frac{1}{2})(2\mu)_n}{(n+\mu)\Gamma(\mu)n!} \rho^{n/2} & \text{if } n \text{ even,} \\ 0 & \text{if } n \text{ odd.} \end{cases} \end{aligned}$$

**Proof.** Same as in the proof of Corollary 7, except apply to both sides of (30). If  $n$  is odd then the Kronecker delta that appears in the orthogonality relation is always equal to zero. ■

**Corollary 10.** *Let  $n \in \mathbf{N}_0$ ,  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ .  
Then*

$$\int_{-1}^1 (1-x^2)^{\mu-1/2} \left( \frac{1}{(1+\rho-2\sqrt{\rho}x)^\mu} - \frac{1}{(1+\rho+2\sqrt{\rho}x)^\mu} \right) C_n^\mu(x) dx$$

$$= \begin{cases} 0 & \text{if } n \text{ even,} \\ \frac{2\sqrt{\pi} \Gamma(\mu + \frac{1}{2})(2\mu)_n}{(n+\mu)\Gamma(\mu)n!} \rho^{n/2} & \text{if } n \text{ odd.} \end{cases}$$

**Proof.** Same as in the proof of Corollary 7, except apply to both sides of (32). If  $n$  is even then the Kronecker delta that appears in the orthogonality relation is always equal to zero. ■

**Corollary 11.** *Let  $n \in \mathbf{N}_0$ ,  $\lambda \in \mathbf{C}$ ,  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ .  
Then*

$$\int_{-1}^1 (1-x^2)^{\mu/2-1/4} P_{\mu-1/2-\lambda}^{1/2-\mu} \left( \sqrt{1+\rho^2-2\rho x} + \rho \right) P_{\mu-1/2-\lambda}^{1/2-\mu} \left( \sqrt{1+\rho^2-2\rho x} - \rho \right) C_n^\mu(x) dx$$

$$= \frac{(\lambda)_n (2\mu-\lambda)_n \rho^{n+\mu-1/2} 2^{\mu-1/2}}{(n+\mu)\Gamma(2\mu)(\mu+1/2)_n n!}.$$

**Proof.** Same as in the proof of Corollary 7, except apply to both sides of (34). ■

**Corollary 12.** *let  $n \in \mathbf{N}_0$ ,  $\alpha \in \mathbf{C}$ ,  $\mu \in (-1/2, \infty) \setminus \{0\}$ ,  $\rho \in (0, 1)$ . Then*

$$\int_{-1}^1 \frac{(1-x^2)^{1/4-\mu/2}}{(1+\rho^2-2\rho x)^{1/4-\mu/2+\alpha/2}} P_{\mu-\alpha-1/2}^{1/2-\mu} \left( \frac{1-\rho x}{\sqrt{1+\rho^2-2\rho x}} \right) C_n^\mu(x) dx$$

$$= \frac{(\alpha)_n \sqrt{\pi} \rho^{n+\mu-1/2} 2^{1/2-\mu}}{(n+\mu)\Gamma(\mu)n!}.$$

**Proof.** Same as in the proof of Corollary 7, except apply to both sides of (35). ■

#### 4. Chebyshev polynomials of the second kind

The Chebyshev polynomials of the second kind can be obtained from the Gegenbauer polynomials using ([?, (18.7.4)]), namely

$$U_n(z) = C_n^1(z), \quad (36)$$

for  $n \in \mathbf{N}_0$ . Hence and through (26), the Chebyshev polynomials of the second kind  $U_n : \mathbf{C} \rightarrow \mathbf{C}$  can be defined in terms of the terminating Gauss hypergeometric series as follows

$$U_n(z) := (n+1) {}_2F_1 \left( \begin{matrix} -n, n+2 \\ \frac{3}{2} \end{matrix} ; \frac{1-z}{2} \right). \quad (37)$$

The orthogonality relation for Chebyshev polynomials of the second kind can be found in ([?, (18.2.1), (18.2.5), Table 18.3.1])

$$\int_{-1}^1 U_m(x) U_n(x) (1-x^2)^{1/2} dx = \frac{\pi}{2} \delta_{m,n}. \quad (38)$$

##### 4.1. Expansions for Chebyshev polynomials of the second kind

**Corollary 13.** *Let  $m \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\} \setminus (-1, 0]$ ,  $x \in [-1, 1]$ . Then*

$$\begin{aligned} & \frac{1}{(1+\rho^2-2\rho x)^{m/2+1}} U_m \left( \frac{1+\rho}{\sqrt{1+\rho^2-2\rho x}} \right) \\ &= \frac{2(m+1)}{\rho(1-\rho)^m} \sum_{n=0}^{\infty} (n+1)(m+2)_{2n} P_{-m}^{-2n-2} \left( \frac{1+\rho}{1-\rho} \right) U_n(x). \end{aligned} \quad (39)$$

**Proof.** Using (28), with (36) and

$$U_m(z) = \sqrt{\frac{\pi}{2}} \frac{m+1}{(z^2-1)^{1/4}} P_{m+1/2}^{-1/2}(z), \quad (40)$$

which follows from (5), (37), and ([?, (15.8.1)]). This completes the proof. ■

Note that using (19), (40) we have the following elementary function representation of the Chebyshev polynomials of the second kind

$$U_m(z) = \frac{1}{2} (z^2-1)^{-1/2} \left[ (z + \sqrt{z^2-1})^{m+1} - (z - \sqrt{z^2-1})^{m+1} \right].$$

**Corollary 14.** *Let  $m \in \mathbf{N}_0$ ,  $\rho \in (0, 1)$ ,  $x \in [-1, 1]$ . Then*

$$\begin{aligned} & \frac{1}{(1 + \rho^2 - 2\rho x)^{m/2+1}} U_m \left( \frac{1 - \rho}{\sqrt{1 + \rho^2 - 2\rho x}} \right) \\ &= \frac{2(m+1)}{\rho(1+\rho)^m} \sum_{n=0}^{\infty} (n+1)(m+2)_{2n} P_{-m}^{-2n-2} \left( \frac{1-\rho}{1+\rho} \right) U_n(x). \end{aligned} \quad (41)$$

**Proof.** Using (29), substitute  $\mu = 1$  with (36) and

$$U_m(x) = \sqrt{\frac{\pi}{2}} \frac{m+1}{(1-x^2)^{1/4}} P_{m+1/2}^{-1/2}(x),$$

for  $x \in (-1, 1)$ , which follows from (?, (14.5.12), (18.5.2)). This completes the proof.  $\blacksquare$

**Corollary 15.** *Let  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in \{z \in \mathbf{C} : z \text{ interior to an ellipse with foci at } \pm 1 \text{ with semi-major axis } (1+\rho)/(2\sqrt{\rho})\}$ . Then*

$$\frac{1}{1 + \rho - 2\sqrt{\rho}x} + \frac{1}{1 + \rho + 2\sqrt{\rho}x} = 2 \sum_{n=0}^{\infty} \rho^n U_{2n}(x). \quad (42)$$

**Proof.** Using (36), and substituting  $\mu = 1$  in (30) with simplification, produces this generating function for Chebyshev polynomials of the second kind.  $\blacksquare$

**Corollary 16.** *Let  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in \{z \in \mathbf{C} : z \text{ interior to an ellipse with foci at } \pm 1 \text{ with semi-major axis } (1+\rho)/(2\sqrt{\rho})\}$ . Then*

$$\frac{1}{1 + \rho - 2\sqrt{\rho}x} - \frac{1}{1 + \rho + 2\sqrt{\rho}x} = 2\sqrt{\rho} \sum_{n=0}^{\infty} \rho^n U_{2n+1}(x). \quad (43)$$

**Proof.** Using (36), and substituting  $\mu = 1$  into (32) with simplification, produces this generating function for Chebyshev polynomials of the second kind.  $\blacksquare$

**Corollary 17.** *Let  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in [-1, 1]$ . Then*

$$(1-x^2)^{-1/4} P_{1/2-\lambda}^{-1/2} \left( \sqrt{1+\rho^2-2\rho x} + \rho \right) P_{1/2-\lambda}^{-1/2} \left( \sqrt{1+\rho^2-2\rho x} - \rho \right) \\ = \frac{2^{5/2} \sqrt{\rho}}{\pi} \sum_{n=0}^{\infty} \frac{(\lambda)_n (2-\lambda)_n 2^{2n} \rho^n}{(2n+2)!} U_n(x). \quad (44)$$

**Proof.** Substituting  $\mu = 1$  into (34), and using (36) with simplification, completes the proof. ■

**Corollary 18.** *Let  $\alpha \in \mathbf{C}$ ,  $\rho \in (0, 1)$ ,  $x \in [-1, 1]$ . Then*

$$\frac{(1+\rho^2-2\rho x)^{1/4-\alpha/2}}{(1-x^2)^{1/4}} P_{1/2-\alpha}^{-1/2} \left( \frac{1-\rho x}{\sqrt{1+\rho^2-2\rho x}} \right) = \sqrt{\frac{2\rho}{\pi}} \sum_{n=0}^{\infty} \frac{(\alpha)_n}{(n+1)!} \rho^n U_n(x). \quad (45)$$

**Proof.** Using (36), and substituting  $\mu = 1$  into (35) with simplification, produces this generating function for Chebyshev polynomials of the second kind. ■

4.2. *Definite integrals for Chebyshev polynomials of the second kind*

**Corollary 19.** *Let  $m, n \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\} \setminus (-1, 0]$ . Then*

$$\int_{-1}^1 \frac{(1-x^2)^{1/2}}{(1+\rho^2-2\rho x)^{m/2+1}} U_m \left( \frac{1+\rho}{\sqrt{1+\rho^2-2\rho x}} \right) U_n(x) dx \\ = \frac{\pi(m+1)(n+1)(m+2)_{2n}}{\rho(1-\rho)^m} P_{-m}^{-2n-2} \left( \frac{1+\rho}{1-\rho} \right).$$

**Proof.** Multiply both sides of (39) by  $(1-x^2)^{1/2} U_k(x)$ , for  $k \in \mathbf{N}_0$  and integrate over  $(-1, 1)$  using the orthogonality relation (38). ■

**Corollary 20.** *Let  $m, n \in \mathbf{N}_0$ ,  $\rho \in (0, 1)$ . Then*

$$\int_{-1}^1 \frac{(1-x^2)^{1/2}}{(1+\rho^2-2\rho x)^{m/2+1}} U_m \left( \frac{1-\rho}{\sqrt{1+\rho^2-2\rho x}} \right) U_n(x) dx \\ = \frac{\pi(m+1)(n+1)(m+2)_{2n}}{\rho(1+\rho)^m} P_{-m}^{-2n-2} \left( \frac{1-\rho}{1+\rho} \right).$$

**Proof.** Same as in the proof of Corollary 19, except apply to both sides of (41). ■

**Corollary 21.** *Let  $n \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ . Then*

$$\int_{-1}^1 (1-x^2)^{1/2} \left( \frac{1}{1+\rho-2\sqrt{\rho}x} + \frac{1}{1+\rho+2\sqrt{\rho}x} \right) U_n(x) dx = \begin{cases} \pi \rho^{n/2} & \text{if } n \text{ even.} \\ 0 & \text{if } n \text{ odd.} \end{cases}$$

**Proof.** Same as in the proof for Corollary 19, except apply to both sides of (42). If  $n$  is odd then the Kronecker delta that appears in the orthogonality relation is always equal to zero. ■

**Corollary 22.** *Let  $n \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ . Then*

$$\int_{-1}^1 (1-x^2)^{1/2} \left( \frac{1}{1+\rho-2\sqrt{\rho}x} - \frac{1}{1+\rho+2\sqrt{\rho}x} \right) U_n(x) dx = \begin{cases} 0 & \text{if } n \text{ even.} \\ \pi \rho^{n/2} & \text{if } n \text{ odd.} \end{cases}$$

**Proof.** Same as in the proof for Corollary 19, except apply to both sides of (43). If  $n$  is even then the Kronecker delta that appears in the orthogonality relation is always equal to zero. ■

**Corollary 23.** *Let  $n \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ . Then*

$$\begin{aligned} \int_{-1}^1 (1-x^2)^{1/4} P_{1/2-\lambda}^{-1/2} \left( \sqrt{1+\rho^2-2\rho x} + \rho \right) P_{1/2-\lambda}^{-1/2} \left( \sqrt{1+\rho^2-2\rho x} - \rho \right) U_n(x) dx \\ = 2^{2n+3/2} \frac{(\lambda)_n (2-\lambda)_n}{(2n+2)!} \rho^{n+1/2}. \end{aligned}$$

**Proof.** Same as in the proof for Corollary 19, except apply to both sides of (44). ■

**Corollary 24.** *Let  $n \in \mathbf{N}_0$ ,  $\alpha \in \mathbf{C}$ ,  $\rho \in (0, 1)$ . Then*

$$\int_{-1}^1 (1-x^2)^{1/4} (1+\rho^2-2\rho x)^{1/4-\alpha/2} P_{1/2-\alpha}^{-1/2} \left( \frac{1-\rho x}{\sqrt{1+\rho^2-2\rho x}} \right) U_n(x) dx = \sqrt{\frac{\pi}{2}} \frac{(\alpha)_n}{(n+1)!} \rho^{n+1/2}.$$

**Proof.** Same as in the proof for Corollary 19, except apply to both sides of (45). ■

## 5. Legendre polynomials

Legendre polynomials can be obtained from the Gegenbauer polynomials using (? , (18.7.9)), namely

$$P_n(z) = C_n^{1/2}(z), \quad (46)$$

for  $n \in \mathbf{N}_0$ . Hence and through (26), the Legendre polynomials of the second kind  $P_n : \mathbf{C} \rightarrow \mathbf{C}$  can be defined in terms of the terminating Gauss hypergeometric series as follows

$$P_n(z) := {}_2F_1 \left( \begin{matrix} -n, n+1 \\ 1 \end{matrix} ; \frac{1-z}{2} \right). \quad (47)$$

The orthogonality relation for Legendre polynomials can be found in (? , (18.2.1), (18.2.5), Table 18.3.1)

$$\int_{-1}^1 P_m(x) P_n(x) dx = \frac{2}{2n+1} \delta_{m,n}. \quad (48)$$

### 5.1. Expansions for Legendre polynomials

**Theorem 10.** *Let  $x, y$  be in a neighborhood of 1. Then*

$$\begin{aligned} & \frac{\pi^2}{2} \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^2}{(n!)^2} P_{2n} \left( \frac{(x+y)(1-xy)}{(x-y)(1+xy)} \right) \left( \frac{x-y}{1+xy} \right)^{2n} \\ &= \begin{cases} \frac{\pi^2}{2} & \text{if } x = y = 1, \\ \frac{1+xy}{xy} K \left( \frac{\sqrt{x^2-1}}{x} \right) K \left( \frac{\sqrt{y^2-1}}{y} \right) & \text{if } x, y \geq 1, \\ \frac{1+xy}{x} K \left( \frac{\sqrt{x^2-1}}{x} \right) K \left( \sqrt{1-y^2} \right) & \text{if } x \geq 1 \text{ and } y \leq 1, \\ \frac{1+xy}{y} K \left( \sqrt{1-x^2} \right) K \left( \frac{\sqrt{y^2-1}}{y} \right) & \text{if } x \leq 1 \text{ and } y \geq 1, \\ (1+xy) K \left( \sqrt{1-x^2} \right) K \left( \sqrt{1-y^2} \right) & \text{if } x, y \leq 1. \end{cases} \end{aligned}$$

**Proof.** If we start with (10) from ?, namely

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n^2}{(n!)^2} P_{2n} \left( \frac{(x+y)(1-xy)}{(x-y)(1+xy)} \right) \left( \frac{x-y}{1+xy} \right)^{2n} \\ = \frac{1+xy}{2} {}_2F_1 \left( \begin{matrix} \frac{1}{2}, \frac{1}{2} \\ 1 \end{matrix} ; 1-x^2 \right) {}_2F_1 \left( \begin{matrix} \frac{1}{2}, \frac{1}{2} \\ 1 \end{matrix} ; 1-y^2 \right), \end{aligned}$$

if we use (?, (15.9.21)) we can express the Gauss hypergeometric functions as Legendre functions. For instance

$${}_2F_1 \left( \begin{matrix} \frac{1}{2}, \frac{1}{2} \\ 1 \end{matrix} ; 1-x^2 \right) = P_{-1/2}(2x^2 - 1),$$

with  $x \in \mathbf{C} \setminus (-\infty, 0]$ . This domain is because the Legendre function of the first kind  $P_\nu$  and the Ferrers function of the first kind  $P_\nu$ , both with order  $\mu = 0$ , are given by the same Gauss hypergeometric function and are continuous across argument unity (cf. (5), (9)). So there is no distinction between these two functions, except that the Ferrers function has argument on the real line with modulus less than unity and the Legendre function is defined on  $\mathbf{C} \setminus (-\infty, 1)$  (both being well defined with argument unity). (Hence there really is no need to use two different symbols to denote this function.) The proof is completed by noting the two formulae

$$P_{-1/2}(z) = \frac{2}{\pi} \sqrt{\frac{2}{z+1}} K \left( \sqrt{\frac{z-1}{z+1}} \right),$$

$$P_{-1/2}(x) = \frac{2}{\pi} K \left( \sqrt{\frac{1-x}{2}} \right)$$

(?, (8.13.1), (8.13.8)), where  $K : [0, 1) \rightarrow [\pi/2, \infty)$  is the complete elliptic integral of the first kind defined by (?, (19.2.8))

$$K(k) := \frac{\pi}{2} {}_2F_1 \left( \begin{matrix} \frac{1}{2}, \frac{1}{2} \\ 1 \end{matrix} ; k^2 \right).$$

■

**Theorem 11.** *Let  $x, y$  be in a neighborhood of 1. Then*

$$= 3 \sum_{n=0}^{\infty} \frac{\left(\frac{1}{3}\right)_n \left(\frac{2}{3}\right)_n}{(n!)^2} P_{3n} \left( \frac{x+y-2x^2y^2}{(x-y)\sqrt{1+4xy(x+y)}} \right) \left( \frac{x-y}{\sqrt{1+4xy(x+y)}} \right)^{3n}.$$

$$= \sqrt{1+4xy(x+y)} \begin{cases} 1 & \text{if } x=y=1 \\ P_{-1/3}(2x^3-1)P_{-1/3}(2y^3-1) & \text{if } x, y > 1 \\ P_{-1/3}(2x^3-1)P_{-1/3}(2y^3-1) & \text{if } x < 1 \text{ and } y > 1 \\ P_{-1/3}(2x^3-1)P_{-1/3}(2y^3-1) & \text{if } x > 1 \text{ and } y < 1 \\ P_{-1/3}(2x^3-1)P_{-1/3}(2y^3-1) & \text{if } x, y < 1 \end{cases}$$

**Proof.** This follows by (?, (11)) and (?, (15.9.21)). ■

**Corollary 25.** *Let  $m \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\} \setminus (-1, 0]$ ,  $x \in [-1, 1]$ . Then*

$$\frac{1}{(1+\rho^2-2\rho x)^{(m+1)/2}} P_m \left( \frac{1+\rho}{\sqrt{1+\rho^2-2\rho x}} \right) = \frac{1}{\sqrt{\rho}(1-\rho)^m} \sum_{n=0}^{\infty} (2n+1)(m+1)_{2n} P_{-m}^{-2n-1} \left( \frac{1+\rho}{1-\rho} \right) P_n(x). \quad (49)$$

**Proof.** Using (28), substitute  $\mu = 1/2$  with (46) and  $P_m(z) = P_m^0(z)$ , which follows from (5), (47). This completes the proof. ■

**Corollary 26.** *Let  $m \in \mathbf{N}_0$ ,  $\rho \in (0, 1)$ ,  $x \in [-1, 1]$ . Then*

$$\frac{1}{(1+\rho^2-2\rho x)^{(m+1)/2}} P_m \left( \frac{1-\rho}{\sqrt{1+\rho^2-2\rho x}} \right) = \frac{1}{\sqrt{\rho}(1+\rho)^m} \sum_{n=0}^{\infty} (2n+1)(m+1)_{2n} P_{-m}^{-2n-1} \left( \frac{1-\rho}{1+\rho} \right) P_n(x). \quad (50)$$

**Proof.** Using (29), substitute  $\mu = 1/2$  with (46) and  $P_m(x) = P_m^0(x)$ , which follows from (9), (47). This completes the proof. ■

**Corollary 27.** *Let  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in \{z \in \mathbf{C} : z \text{ interior to an ellipse with foci at } \pm 1 \text{ with semi-major axis } (1 + \rho)/(2\sqrt{\rho})\}$ . Then*

$$\frac{1}{\sqrt{1 + \rho - 2\sqrt{\rho}x}} + \frac{1}{\sqrt{1 + \rho + 2\sqrt{\rho}x}} = 2 \sum_{n=0}^{\infty} \rho^n P_{2n}(x). \quad (51)$$

**Proof.** Substituting  $\mu = 1/2$  in (30) and using (46) with simplification, produces this generating function for Legendre polynomials. ■

**Corollary 28.** *Let  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in \{z \in \mathbf{C} : z \text{ interior to an ellipse with foci at } \pm 1 \text{ with semi-major axis } (1 + \rho)/(2\sqrt{\rho})\}$ . Then*

$$\frac{1}{\sqrt{1 + \rho - 2\sqrt{\rho}x}} - \frac{1}{\sqrt{1 + \rho + 2\sqrt{\rho}x}} = 2\sqrt{\rho} \sum_{n=0}^{\infty} \rho^n P_{2n+1}(x). \quad (52)$$

**Proof.** Substituting  $\mu = 1/2$  in (32) and using (46) with simplification, produces this generating function for Legendre polynomials. ■

**Corollary 29.** *Let  $\lambda \in \mathbf{C}$ ,  $\rho \in \{x \in \mathbf{C} : |x| < 1\}$ ,  $x \in [-1, 1]$ . Then*

$$P_{-\lambda} \left( \sqrt{1 + \rho^2 - 2\rho x} + \rho \right) P_{-\lambda} \left( \sqrt{1 + \rho^2 - 2\rho x} - \rho \right) = \sum_{n=0}^{\infty} \frac{(\lambda)_n (1 - \lambda)_n}{(n!)^2} \rho^n P_n(x). \quad (53)$$

**Proof.** Substituting  $\mu = 1/2$  into (34) and using (46) with simplification completes the proof. ■

Note that Corollary 29 is just a restatement of (? , Theorem A), and therefore (8) is a generalization of Brafman's theorem.

**Corollary 30.** *Let  $\alpha \in \mathbf{C}$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ ,  $x \in [-1, 1]$ . Then*

$$(1 + \rho^2 - 2\rho x)^{-\alpha/2} P_{\alpha-1} \left( \frac{1 - \rho x}{\sqrt{1 + \rho^2 - 2\rho x}} \right) = \sum_{n=0}^{\infty} \frac{(\alpha)_n}{n!} \rho^n P_n(x). \quad (54)$$

**Proof.** Substituting  $\mu = 1/2$  in (35) with simplification completes the proof. ■

### 5.2. Definite integrals for Legendre polynomials

**Corollary 31.** *Let  $m, n \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\} \setminus (-1, 0]$ . Then*

$$\int_{-1}^1 \frac{1}{(1 + \rho^2 - 2\rho x)^{(m+1)/2}} P_m \left( \frac{1 + \rho}{\sqrt{1 + \rho^2 - 2\rho x}} \right) P_n(x) dx = \frac{2(m+1)_{2n}}{\sqrt{\rho}(1 - \rho)^m} P_{-m}^{-2n-1} \left( \frac{1 + \rho}{1 - \rho} \right).$$

**Proof.** Multiply both sides of (49) by  $P_k(x)$ , for  $k \in \mathbf{N}_0$  and integrate over  $(-1, 1)$  using the orthogonality relation (48). ■

**Corollary 32.** *Let  $m, n \in \mathbf{N}_0$ ,  $\rho \in (0, 1)$ . Then*

$$\int_{-1}^1 \frac{1}{(1 + \rho^2 - 2\rho x)^{(m+1)/2}} P_m \left( \frac{1 - \rho}{\sqrt{1 + \rho^2 - 2\rho x}} \right) P_n(x) dx = \frac{2(m+1)_{2n}}{\sqrt{\rho}(1 + \rho)^m} P_{-m}^{-2n-1} \left( \frac{1 - \rho}{1 + \rho} \right).$$

**Proof.** Same as in the proof of Corollary 31, except apply to both sides of (50). ■

**Corollary 33.** *Let  $n \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ . Then*

$$\int_{-1}^1 \left( \frac{1}{\sqrt{1 + \rho - 2\sqrt{\rho}x}} + \frac{1}{\sqrt{1 + \rho + 2\sqrt{\rho}x}} \right) P_n(x) dx = \begin{cases} \frac{4\rho^{n/2}}{2n+1} & \text{if } n \text{ even,} \\ 0 & \text{if } n \text{ odd.} \end{cases}$$

**Proof.** Same as in the proof of Corollary 31, except apply to both sides of (51). If  $n$  is odd then the Kronecker delta that appears in the orthogonality relation is always equal to zero. ■

**Corollary 34.** *Let  $n \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ . Then*

$$\int_{-1}^1 \left( \frac{1}{\sqrt{1 + \rho - 2\sqrt{\rho}x}} - \frac{1}{\sqrt{1 + \rho + 2\sqrt{\rho}x}} \right) P_n(x) dx = \begin{cases} 0 & \text{if } n \text{ even.} \\ \frac{4\rho^{n/2}}{2n+1} & \text{if } n \text{ odd.} \end{cases}$$

**Proof.** Same as in the proof of Corollary 31, except apply to both sides of (52). If  $n$  is even then the Kronecker delta that appears in the orthogonality relation is always equal to zero. ■

**Corollary 35.** *Let  $n \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ . Then*

$$\int_{-1}^1 P_\lambda \left( \sqrt{1 + \rho^2 - 2\rho x + \rho} \right) P_\lambda \left( \sqrt{1 + \rho^2 - 2\rho x - \rho} \right) P_n(x) dx = \frac{2(\lambda + 1)_n (-\lambda)_n}{(2n + 1)(n!)^2} \rho^n.$$

**Proof.** Same as in the proof of Corollary 31, except apply to both side of (53). ■

**Corollary 36.** *Let  $n \in \mathbf{N}_0$ ,  $\alpha \in \mathbf{C}$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\}$ . Then*

$$\int_{-1}^1 (1 + \rho^2 - 2\rho x)^{-\alpha/2} P_{\alpha-1} \left( \frac{1 - \rho x}{\sqrt{1 + \rho^2 - 2\rho x}} \right) P_n(x) dx = \frac{2(\alpha)_n \rho^n}{(2n + 1)n!}.$$

**Proof.** Same as in the proof of Corollary 31, except apply to both sides of (54). ■

## 6. Chebyshev polynomials of the first kind

The Chebyshev polynomials of the first kind  $T_n : \mathbf{C} \rightarrow \mathbf{C}$  can be defined in terms of the terminating Gauss hypergeometric series as follows ([?, p. 257])

$$T_n(z) := {}_2F_1 \left( \begin{matrix} -n, n \\ \frac{1}{2} \end{matrix} ; \frac{1 - z}{2} \right), \quad (55)$$

for  $n \in \mathbf{N}_0$ . The Chebyshev polynomials of the first kind can be obtained from the Gegenbauer polynomials using ([?, (6.4.13)]), namely

$$T_n(z) = \frac{1}{\epsilon_n} \lim_{\mu \rightarrow 0} \frac{n + \mu}{\mu} C_n^\mu(z), \quad (56)$$

where the Neumann factor  $\epsilon_n \in \{1, 2\}$ , commonly seen in Fourier cosine series, is defined as  $\epsilon_n := 2 - \delta_{n,0}$ . The orthogonality relation for Chebyshev polynomials of the first kind can be found in ([?, (18.2.1), (18.2.5), Table 18.3.1])

$$\int_{-1}^1 T_m(x) T_n(x) (1 - x^2)^{-1/2} dx = \frac{\pi}{\epsilon_n} \delta_{m,n}. \quad (57)$$

6.1. Expansions for Chebyshev polynomials of the first kind

**Corollary 37.** *Let  $m \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\} \setminus (-1, 0]$ ,  $x \in [-1, 1]$ . Then*

$$\frac{1}{(1 + \rho^2 - 2\rho x)^{m/2}} T_m \left( \frac{1 + \rho}{\sqrt{1 + \rho^2 - 2\rho x}} \right) = \frac{1}{(1 - \rho)^m} \sum_{n=0}^{\infty} \epsilon_n(m)_{2n} P_{-m}^{-2n} \left( \frac{1 + \rho}{1 - \rho} \right) T_n(x). \quad (58)$$

**Proof.** Using (28), (56), and

$$T_m(z) = \sqrt{\frac{\pi}{2}} (z^2 - 1)^{1/4} P_{m-1/2}^{1/2}(z), \quad (59)$$

which follows from (5), (55), (?, (15.8.1)). This completes the proof.  $\blacksquare$

Note that using (18), (59) we have the following elementary function representation of the Chebyshev polynomials of the first kind (see also (?, p. 177))

$$T_m(z) = \frac{1}{2} \left[ (z + \sqrt{z^2 - 1})^m - (z - \sqrt{z^2 - 1})^m \right].$$

**Corollary 38.** *Let  $m \in \mathbf{N}_0$ ,  $\rho \in (0, 1)$ ,  $x \in [-1, 1]$ . Then*

$$\frac{1}{(1 + \rho^2 - 2\rho x)^{m/2}} T_m \left( \frac{1 - \rho}{\sqrt{1 + \rho^2 - 2\rho x}} \right) = \frac{1}{(1 + \rho)^m} \sum_{n=0}^{\infty} \epsilon_n(m)_{2n} P_{-m}^{-2n} \left( \frac{1 - \rho}{1 + \rho} \right) T_n(x). \quad (60)$$

**Proof.** Using (29), (56), and

$$T_m(x) = \sqrt{\frac{\pi}{2}} (1 - x^2)^{1/4} P_{m-1/2}^{1/2}(x)$$

which follows from (9), (55), (?, (15.8.1)). This completes the proof.  $\blacksquare$

6.2. Definite integrals for Chebyshev polynomials of the first kind

**Corollary 39.** Let  $m, n \in \mathbf{N}_0$ ,  $\rho \in \{z \in \mathbf{C} : |z| < 1\} \setminus (-1, 0]$ . Then

$$\int_{-1}^1 \frac{(1-x^2)^{-1/2}}{(1+\rho^2-2\rho x)^{m/2}} T_m \left( \frac{1+\rho}{\sqrt{1+\rho^2-2\rho x}} \right) T_n(x) dx = \frac{2\pi(m)_{2n}}{\epsilon_n(1-\rho)^m} P_{-m}^{-2n} \left( \frac{1+\rho}{1-\rho} \right).$$

**Proof.** Multiply both sides of (58) by  $(1-x^2)^{-1/2} T_k(x)$ , for  $k \in \mathbf{N}_0$  and integrate over  $(-1, 1)$  using the orthogonality relation (57). ■

**Corollary 40.** Let  $m, n \in \mathbf{N}_0$ ,  $\rho \in (0, 1)$ . Then

$$\int_{-1}^1 \frac{(1-x^2)^{-1/2}}{(1+\rho^2-2\rho x)^{m/2}} T_m \left( \frac{1-\rho}{\sqrt{1+\rho^2-2\rho x}} \right) T_n(x) dx = \frac{2\pi(m)_{2n}}{\epsilon_n(1+\rho)^m} P_{-m}^{-2n} \left( \frac{1-\rho}{1+\rho} \right).$$

**Proof.** Same as in the proof of Corollary 39 except apply to both sides of (60). ■

*Acknowledgements*

The authors thank Hans Volkmer for valuable discussions. C. MacKenzie would like to thank the Summer Undergraduate Research Fellowship program at the National Institute of Standards and Technology for financial support while this research was carried out. This work was conducted while H. S. Cohl was a National Research Council Research Postdoctoral Associate in the Information Technology Laboratory at the National Institute of Standards and Technology, Gaithersburg, Maryland, U.S.A.